

Spring 5-3-2018

Institutional Outcomes following Total Pancreatectomy and Islet Autotransplantation in Children

Stephen P. Sekoulopoulos
Valparaiso University, stephen.sekoulopoulos@valpo.edu

Follow this and additional works at: <https://scholar.valpo.edu/cus>

Recommended Citation

Sekoulopoulos, Stephen P., "Institutional Outcomes following Total Pancreatectomy and Islet Autotransplantation in Children" (2018). *Symposium on Undergraduate Research and Creative Expression (SOURCE)*. 731.
<https://scholar.valpo.edu/cus/731>

This Poster Presentation is brought to you for free and open access by the Office of Sponsored and Undergraduate Research at ValpoScholar. It has been accepted for inclusion in Symposium on Undergraduate Research and Creative Expression (SOURCE) by an authorized administrator of ValpoScholar. For more information, please contact a ValpoScholar staff member at scholar@valpo.edu.

Institutional Outcomes following Total Pancreatectomy with Islet Autotransplantation in Children

Stephen Sekoulopoulos and Dr. Jaimie Nathan

Cincinnati Children's Hospital Division of Pediatric General and Thoracic Surgery

Aim

- To review our institutional outcomes following total pancreatectomy and islet autotransplantation for debilitating acute recurrent (ARP) and chronic pancreatitis (CP) in children.

Background

- Chronic pancreatitis is a progressive fibroinflammatory injury that can result in endocrine and/or exocrine insufficiency, as well as in severe pain and impaired quality of life.
- The fibrotic pancreas is removed while the insulin-producing islets are isolated and transplanted back into the patient, usually into the liver via the portal vein.

Figure 1: Total pancreatectomy






Figure 2: Biliary and Alimentary Tract Reconstruction

Figure 3: Islet Autotransplantation



Methodology

- 17 patients underwent pancreatectomy with islet autotransplantation between April 2015 and April 2017 at Cincinnati Children's Hospital.
- Open, pylorus-preserving approach with accompanied splenectomy and gastrojejunostomy tube placement.
- 16 total pancreatectomies and 1 subtotal pancreatectomy.
- Islet isolation by enzymatic and mechanical digestion (Ricordi method).
- Documented surgical complications, readmission, reoperations 90 days post-op.
- Preoperative and postoperative opioid use was assessed, as was islet function.

Results

Table 1: Demographic Data

Criteria	Values (N = 17)
Sex (female)	11 (65%)
Age (years)	12.1 (4 - 17)
Weight (kg)	53.6 (16.6 – 108.3)
Height (cm)	150.2 (103 – 185.5)
Acute Recurrent Pancreatitis (yes)	3 (18%)
Chronic Pancreatitis (yes)	14 (82%)
Preoperative Enteral Tube or TPN (#)	7 (41%)
Visits for Pain 18 months prior to operation (#)	2.5 (0 – 6)
ED visits 18 months prior to operation (#)	1.3 (0 – 6)
Genetic mutations (yes)	14 (82%)
PRSS1	6 (35%)
SPINK1	5 (29%)
CFTR	6 (35%)
CTRC	0 (0%)
Pancreas Divisum (yes)	8 (47%)
Autoimmune Pancreatitis (yes)	1 (6%)
Pancreatic Exocrine Insufficiency (yes)	8 (47%)
Required Pre-Operative Insulin	1 (6%)
Postop ICU Length of Stay (days)	10.4 (7 – 18)
Total Length of Hospital Stay (days)	26.1 (19 – 36)





Table 2: Post-Operative Complications

Complication	Values (N = 17)
Intra-Abdominal Bleeding	1 (6%)
Upper GI Bleeding	1 (6%)
Small Bowel Obstruction	1 (6%)
Wound Infection	2 (12%)
Portal Vein Thrombosis	0 (0%)
Fluid Collection	3 (18%)
Hypoglycemia	1 (6%)

Table 3: Readmission and Reoperation following TPIAT

Criteria	Values (N = 17)
Number of Patients Readmitted	6 (35%)
Fever	1 (6%)
Abdominal or Other Non-Localized Pain	4 (24%)
Enteral Tube Complications	2 (12%)
Length of Readmission (days)	4.9 (2 - 12)
Patients Returning to the OR	2 (12%)
Exploratory Laparotomy	2 (12%)
Patients Requiring an IR/Endo Procedure	6 (35%)
Drainage of Intra-abdominal Fluid	3 (18%)
PICC	2 (12%)
C-Line Removal	2 (12%)
EGD	1 (6%)
GJ Tube Change	2 (12%)
Therapeutic Procedure	2 (12%)

Table 4: Islet Infusion and Insulin Use

Criteria	Values (N = 17)
Total IEQ	330,326 (172,300 – 488,000)
IEQ/kg	6,677 (3,992 – 14,242)
Portal Vein Infusion	16 (94%)
Peritoneal Cavity Infusion	1 (6%)
Insulin Independent	4 (24%)
Time on Insulin (months)	8 (4 – 16)
Partial Graft Function	13 (76%)

Conclusions

- TPIAT is an important option in treatment of children with ARP/CP and is considered when maximal medical therapies and endoscopic approaches fail to relieve pain and to address complications.
- Comprehensive multidisciplinary patient evaluation is critical to ensure optimal outcomes following TPIAT.
- In appropriately selected children, TPIAT achieves durable pain relief and improves QOL with manageable glycemic control.